FAT INJECTION

DR. GERUT'S POST OPERATIVE INSTRUCTIONS

Avoid, talking, chewing, etc. Do not use straws. DO NOT CHEW GUM OR SMOKE!

On the first night after surgery, avoid excessive chewing. After the second day you may eat or drink anything that you wish.

Please take the medications that Dr. Gerut indicates with a check mark. Only the medications checked off will be given to you. If you are allergic to any of these medications, please let us know! Please take all pills with a bit of food to avoid nausea. When you fill your prescriptions, the bottles will be labeled only with the medical names for your pills and for your reference, these medical names are in parentheses below.

Antibiotic:	□ Keflex (Cephalexin) 500mg (1pill) 2 times a day Or, if allergic to Penicillin,		
	□Vibramycin (Doxycycline) - 100mg (1pill) 2 times a day		
Severe pain:	□Percocet (Oxycodone) - 1-2 pills every 3 to 4 hours as needed.		
Moderate pain:	■ Norco (hydrocodone bitartrate/acetaminophen)- 1-2 pills every 3 to 4 hours as needed.		
Nausea:	☐ Compazine (Prochloperazine) Use one if needed. Never use more than 2 in 24 hours. If nausea returns, and you weigh less than 160 pounds, call the office before using a second suppository.		
Constipation:	Ex-Lax (or any other mild laxative as needed).		
Multi-Vitamin:	□1 daily		

There will be a certain amount of swelling that will last a variable number of days. To minimize swelling, keep your head elevated at all times. When lying down or sleeping, sleep in a reclining chair or use as may pillows as possible. You may apply ice if you wish. If you have a any problem especially a sudden increase in pain or a fever (measured by a thermometer) or if you have MUCH more swelling on one side compared to the other, **call the office at 516-295-2100**. After office hours please call **516-620-3619**.

I have been made aware of the fol them (there may be other problem		ns and/or side effects and I accept
Bleeding Need for revisional surgery Change in facial shape Parasthesias and/or numbness Change in facial expression X	Reaction to the sutures Chronic pain Seroma Hematoma	Infection Asymmetry Anesthesia risks Contour deformities
It may be necessary for you to consurgery. This communication may over the Internet from which you acknowledge and accept this pote	y involve sending informatio may or may not be identifiab	n about or photographs of yourself
No Aspirin, Bufferin, APC, Fiorin other medication not directed by I Advil or Motrin, check with Dr. Conarcotic pain reliever, you are to a severe drowsiness can cause falls	Or. Gerut for ONE WEEK Algerut or his office staff. During or operating heads	FTER SURGERY. Regarding ng the time that you are taking a
Significant weight gain or weight surgical result, For this reason, you after your procedure. If your weight) after your procedure, you same area and you will be charged fee. If for other reasons, revisional there is no surgical fee, THERE VIROOM AND THE ANESTHESIA	ou must agree to make all efforght changes significantly (changes will not be entitled to any frod a surgical fee and an operatal surgery is required, there moved the surgery is required.	orts to keep your weight stable ange of 10% or more in body see "touch-up" procedures to the ing room fee and an anesthesia ay be a surgical fee but even if
I have read the above instructions have been answered and I underst	, , , ,	
Patient Signature	I	Date
Zachary E. Gerut, M.D., F.A.C.S., P.C. Diplomat, American Board of Plastic Su Member, American Society of Plastic Su Member, American Society for Aesthetic Assistant Clinical Professor of Plastic Su Medical Director or Research, Touro Un Fellow of the American College of Surg	orgeons o Plastic Surgery orgery, Albert Einstein Med Ctr. (re oriversity Physician Assistant Schoo	
or and rametream comege of built		10, 0, 10